


LADYBIRD BEETLES OF IDAHO

Ladybird beetles (Coccinellidae) are significant predators of aphids, mealybugs and scale insects and can also prey on other small arthropods. A few groups of species feed on fungi or plant tissue. Eighty-eight species are known in Idaho, all but a few uncommon species illustrated here, with some color variants. Distribution maps are based on museum specimen labels (Barr Museum in Moscow, Orma Smith Museum in Caldwell) and online biodiversity databases (<https://www.gbif.org/>).


COCCINELLINAE




Adalia bipunctata
Twospotted lady beetle
(3.5-5.2 mm)




Anatis lecontei
(7.8-10.5 mm)




Anatis mali
(7.3-10.0 mm)




Anatis rathvoni
(7.5-10.2 mm)




Anisosticta bitriangularis
(3.0-4.0 mm)





Calvia quatuordecimguttata
(4.0-5.5 mm)




Coccinella difficilis
(5.0-6.0 mm)



Coccinella monticola
(5.2-7.0 mm)




Coccinella novemnotata
Ninespotted lady beetle
(4.7-7.0 mm)




Coccinella prolongata
(5.7-7.8 mm)




Coccinella septempunctata
Sevenspotted lady beetle
(6.5-7.8 mm)



Coccinella transversoguttata
Transverse lady beetle
(5.0-7.8 mm)



Coccinella trifasciata
(4.0-5.0 mm)



Cycloneda polita
(3.5-6.2 mm)



Harmonia axyridis
Multicolored Asian lady beetle
(5.5-8.5 mm)




Hippodamia apicalis
(3.5-4.7 mm)




Hippodamia caseyi
(4.8-6.7 mm)




Hippodamia convergens
Convergent lady beetle
(4.2-7.3 mm)




Hippodamia expurgata
(3.5-5.0 mm)




Hippodamia falcigera
(5.0-6.0 mm)



Hippodamia glacialis
(5.0-7.0 mm)




Hippodamia moesta
(6.0-7.5 mm)



Hippodamia parenthesis
(3.8-5.6 mm)




Hippodamia quinquesignata
(4.0-7.0 mm)




Hippodamia sinuata
Sinuate lady beetle
(4.3-5.8 mm)





Hippodamia tredecimpunctata
Thirteenspotted lady beetle
(4.5-6.4 mm)




Hippodamia washingtoni
(5.4-6.7 mm)




Macronaemia episcopalis
(3.3-4.0 mm)




Mulsantina picta
(3.3-5.3 mm)





Myzia interrupta
(6.5-8.0 mm)





Myzia subvittata
(5.7-8.0 mm)




Paraneamia vittigera
(4.5-6.6 mm)




Psyllobora borealis
(2.4-3.1 mm)





Psyllobora vigintimaculata
(1.8-3.0 mm)




Epilachna varivestis
Mexican bean beetle
(6.4-8.1 mm)



Microweisa misella
(1.0-1.5 mm)



Brumoides septentrionis
(2.9-3.0 mm)




Chilocorus stigma
Twicestabbed lady beetle
(3.8-5.0 mm)



Exochomus aethiops
(3.0-4.2 mm)

SCYMNINAE




Brachiantha utelela
(2.6-4.6 mm)




Hyperaspis hercules
(2.1-4.0 mm)




Hyperaspis mimus
(1.4-1.8 mm)




Hyperaspis vittigerus
(1.5-2.1 mm)





Hyperaspis chapini
(2.0-2.8 mm)





Hyperaspis dissoluta
(2.0-3.1 mm)





Hyperaspis fastidiosa
(2.1-2.8 mm)



Hyperaspis inflexa
(1.8-2.9 mm)




Hyperaspis lateralis
(2.5-3.8 mm)




Hyperaspis oregona
(2.0-2.6 mm)





Hyperaspis pleuralis
(2.0-2.9 mm)




Hyperaspis postica
(2.3-3.1 mm)



Hyperaspis quadrivittata
(2.0-2.7 mm)



Hyperaspis undulata
(1.8-2.8 mm)



Nephus atramentarius
(1.7-2.1 mm)




Nephus ornatus
(1.7-2.1 mm)




Nephus sordidus
Little brown mealybug destroyer
(1.4-1.8 mm)




Scymnus calaveras
(2.1-2.8 mm)




Scymnus caurinus
(1.8-2.4 mm)




Scymnus coniferarum
(1.6-2.1 mm)





Scymnus fenderi
(2.2-2.6 mm)



Scymnus humboldti
(2.0-2.6 mm)



Scymnus jacobianus
(1.6-2.3 mm)




Scymnus lacustris
(1.1-1.5 mm)



Scymnus loewii
(1.8-2.4 mm)


COCCIDULINAE



Scymnus marginicollis
(1.8-2.2 mm)



Scymnus nebulosus
(2.0-2.5 mm)




Scymnus postoictus
(1.7-2.2 mm)



Scymnus solidus
(2.0-2.6 mm)





Scymnus tahoensis
(1.1-1.5 mm)



Stethorus punctum
Spider mite destroyer
(1.4-1.6 mm)



Zilus aterrimus
Spider mite destroyer
(1.3-1.6 mm)



Coccidula lepida
(2.8-3.5 mm)

- Feeding habits:**
- Aphids
 - Mealybugs/scales
 - Spider mites
 - Fungus (mildew)
 - Bean plant leaves, flowers and pods

© Copyright 2025 University of Idaho Barr Entomological Museum. Downloading, printing, sharing and distribution are allowed. Partial reproduction for non-commercial purposes allowed if original source is acknowledged. Unauthorized sale is strictly prohibited. This poster was compiled by Eli Carmona (photography, photo editing, distribution data) and Luc Leblanc (research, design, maps). All photographed specimens are digitized (<https://ecdysis.org>) and deposited at the William F. Barr Entomological Museum (University of Idaho, Moscow, ID). Photos of *Hippodamia falcigera* and *H. washingtoni* by Don Griffiths, courtesy of the Spencer Entomological Collection, Beaty Biodiversity Museum, UBC.